

Effectiveness of Yoga.docx

 My Files

 My Files

 NUSRL, Ranchi

Document Details

Submission ID

trn:oid::3618:133391456

Submission Date

Mar 29, 2026, 4:31 PM GMT+5:30

Download Date

Mar 29, 2026, 4:35 PM GMT+5:30

File Name

Effectiveness of Yoga.docx

File Size

172.2 KB

13 Pages

5,589 Words

41,540 Characters

9% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Filtered from the Report

- ▶ Bibliography
- ▶ Quoted Text
- ▶ Cited Text
- ▶ Small Matches (less than 8 words)

Match Groups

- 56 Not Cited or Quoted 9%**
 Matches with neither in-text citation nor quotation marks
- 0 Missing Quotations 0%**
 Matches that are still very similar to source material
- 0 Missing Citation 0%**
 Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**
 Matches with in-text citation present, but no quotation marks

Top Sources

- 7% Internet sources
- 3% Publications
- 5% Submitted works (Student Papers)

Integrity Flags

1 Integrity Flag for Review

- Hidden Text**
 5585 suspect characters on 13 pages
 Text is altered to blend into the white background of the document.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- 56 Not Cited or Quoted 9%**
Matches with neither in-text citation nor quotation marks
- 0 Missing Quotations 0%**
Matches that are still very similar to source material
- 0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 7% Internet sources
- 3% Publications
- 5% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Publication	Yaser Mohammed Al-Worafi. "Handbook of Complementary, Alternative, and Inte...	<1%
2	Internet	www.mdpi.com	<1%
3	Submitted works	Nims University Rajasthan, Jaipur on 2026-02-04	<1%
4	Submitted works	Manipal University Jaipur Online on 2026-01-12	<1%
5	Internet	link.springer.com	<1%
6	Internet	ijmrr.medresearch.in	<1%
7	Internet	getmarlee.com	<1%
8	Submitted works	MAHSA UNIVERSITY (NEW LMS) on 2026-03-26	<1%
9	Internet	vestnik-philological.tou.edu.kz	<1%
10	Internet	www.academypublication.com	<1%

11	Submitted works	South Cheshire on 2026-03-27	<1%
12	Internet	scholar.archive.org	<1%
13	Internet	healthdor.com	<1%
14	Internet	www.kheljournal.com	<1%
15	Submitted works	Concordia University Irvine on 2026-02-09	<1%
16	Submitted works	Curtin University LTI on 2026-03-16	<1%
17	Internet	admin.mantechpublications.com	<1%
18	Internet	www.frontiersin.org	<1%
19	Submitted works	Atharva College of Engineering on 2026-03-27	<1%
20	Internet	edoc.hu-berlin.de	<1%
21	Internet	www.bursahaga.com	<1%
22	Internet	fishtaxa.com	<1%
23	Internet	www.allstudyjournal.com	<1%
24	Internet	www.stevenhuff.net	<1%

25	Publication	Denham-Jones, Laura. "A Comparison of the Effects of Yoga and Pilates for Pain, P..."	<1%
26	Publication	Garcia, Gryan Eroll Robles. "Clinical Correlation Between Hematological Indices a..."	<1%
27	Submitted works	Glasgow Caledonian University on 2024-08-12	<1%
28	Submitted works	Institute of Graduate Studies, UiTM on 2014-06-27	<1%
29	Submitted works	Mt San Antonio College on 2025-11-22	<1%
30	Publication	Siddharth Sagle, Neetu Kumari, Priya Bhardwaj, Mulaka Maruthi, Ashok Jangra, D...	<1%
31	Publication	Susan Holtzman, R Thomas Beggs. "Yoga for Chronic Low Back Pain: A Meta-Anal..."	<1%
32	Internet	bioclima.ro	<1%
33	Submitted works	hcpc on 2025-08-08	<1%
34	Internet	ijsrem.com	<1%
35	Internet	inspiree.review	<1%
36	Internet	journalijsra.com	<1%
37	Internet	mdpi-res.com	<1%
38	Internet	mlrb.net	<1%

39	Internet	periodicos.feevale.br	<1%
40	Internet	pubs.sciepub.com	<1%
41	Internet	repository.cpu.edu.ph	<1%
42	Internet	rsisinternational.org	<1%
43	Internet	wiredspace.wits.ac.za	<1%
44	Internet	www.hnjournal.net	<1%
45	Internet	www.versalie.com	<1%

“Effectiveness of Yoga-Based Balance Exercises versus Conventional Balance Training on Functional Mobility and Quality of Life among Elderly Individuals

Abstract

The present study used a comparative experimental research design to evaluate the effectiveness of yoga-based balance exercises and conventional balance training on functional mobility and quality of life among elderly individuals aged 60 years and above. A total of 40 participants were selected using purposive sampling and randomly allocated into two intervention groups with 20 participants in each group. The yoga group performed structured yoga-based balance exercises while the comparison group performed conventional physiotherapy-based balance training. Both intervention programs were conducted for 8 weeks three sessions per week with each session lasting approximately 45 minutes under the supervision of a trained instructor. Functional mobility was assessed using standardized measures including the Timed Up and Go (TUG) test and gait speed test while quality of life was evaluated using the WHOQOL-BREF questionnaire. Pre-test and post-test assessments were conducted to evaluate the effectiveness of the interventions. The findings indicated improvements in functional mobility and quality of life in both groups; however the yoga-based balance exercise group demonstrated relatively greater improvement in overall quality-of-life scores. These findings suggest that yoga-based exercises may provide additional benefits as a complementary intervention for improving mobility and well-being among elderly individuals.

Keywords: Balance Training Functional Mobility Comparative Experimental Complementary Alternative Experimental Research

1. Introduction

Background of Aging and Decline in Balance

Aging is a natural biological process characterized by progressive physiological psychological and functional changes that affect the overall health and independence of individuals. One of the most significant changes associated with aging is the gradual decline in balance and postural control. Balance is a complex physiological function that involves the integration of sensory input from the visual vestibular and proprioceptive systems along with appropriate neuromuscular responses to maintain stability during both static and dynamic activities. As individuals age these systems gradually deteriorate. The vestibular system becomes less sensitive visual acuity declines and proprioceptive feedback from joints and muscles becomes less accurate. In addition age-related reductions in muscle strength flexibility and reaction time further compromise postural stability. These physiological changes contribute to impaired balance and increase the likelihood of instability during everyday activities such as walking standing or climbing stairs.

Prevalence Consequences of Falls,among,the,Elderly

Falls represent one of the most common and serious health concerns among older adults worldwide. According to global health reports, approximately one-third of individuals aged 65 years and above experience at least one fall each year. The incidence of falls tends to increase with advancing age due to declining physical capabilities, medical conditions and environmental hazards. Falls among elderly individuals can lead to severe physical injuries, including fractures, head trauma and soft tissue injuries. Hip fractures, in particular, are among the most serious consequences of falls and are associated with prolonged hospitalization, disability and increased mortality rates. Beyond physical injuries, falls often have significant psychological consequences. Many elderly individuals develop a fear of falling after experiencing a fall, which may lead to reduced physical activity, social isolation

and decreased confidence in performing daily tasks. Furthermore, falls place a substantial burden on healthcare systems and caregivers. They often require medical treatment, rehabilitation and long-term care, which increases healthcare costs and reduces the overall quality of life for older adults. Preventing falls has therefore become a major public health priority and researchers have emphasized the importance of interventions that improve balance, strength and functional mobility in elderly populations.¹

Importance of Balance Training

Balance training is widely recognized as an effective intervention for improving postural stability and reducing the risk of falls among older adults. It involves a variety of exercises designed to enhance coordination, strength, flexibility and sensory integration required for maintaining body equilibrium. Regular balance training can improve neuromuscular control, strengthen lower limb muscles and enhance the body's ability to respond to external disturbances. Several studies have demonstrated that structured balance training programs can significantly improve functional mobility, gait stability and confidence in performing daily activities among elderly individuals. Exercises such as weight shifting, single-leg standing, dynamic gait training and resistance exercises are commonly included in rehabilitation programs aimed at fall prevention. These exercises help improve muscle strength and joint stability, which are essential for maintaining balance during movement. In addition to physical benefits, balance training may also contribute to improved psychological well-being by increasing self-confidence and reducing fear of falling. As a result, elderly individuals become more active and engaged in daily activities, which further supports their overall health and independence. For these reasons, balance training is considered a key component of geriatric rehabilitation and preventive healthcare programs.

Role of Yoga in Geriatric Health

Yoga is an ancient mind-body practice that originated in India and has gained global recognition for its health benefits. It combines physical postures (asanas), breathing techniques (pranayama) and meditation or relaxation practices to promote physical, mental and emotional well-being. In recent years, yoga has been increasingly incorporated into health promotion and rehabilitation programs for elderly populations. Yoga-based exercises can help improve flexibility, muscle strength, balance and coordination, which are essential for maintaining functional independence among older adults. Many yoga postures require controlled movements and sustained body alignment, which can enhance postural awareness and stability. Additionally, breathing exercises and relaxation techniques may reduce stress, improve concentration and promote mental clarity. Research has shown that regular yoga practice can lead to improvements in balance, mobility and physical performance in elderly individuals. It may also help reduce symptoms of anxiety, depression and chronic pain while enhancing overall quality of life.³ Unlike conventional exercise programs that focus primarily on physical fitness, yoga adopts a holistic approach that addresses both physical and psychological aspects of health. This integrated approach makes yoga a promising intervention for improving the well-being of older adults.

Need for Comparative Studies between Yoga and Conventional Balance Training

¹Gillespie L.D. et al. Interventions for preventing falls in older people. *Cochrane Database of Systematic Reviews*.

²Howe T.E. et al. Exercise for improving balance in older people. *Cochrane Review*.

³Tiedemann A. et al. The effect of yoga on balance and mobility in older adults. *Age and Ageing*.

Although both yoga-based exercises and conventional balance training have demonstrated benefits for improving mobility and stability in elderly individuals, there is still a need for more comparative research examining their relative effectiveness. Conventional balance training programs used in physiotherapy typically emphasize strengthening, coordination and functional mobility exercises designed to prevent falls and improve physical performance.⁴ In contrast, yoga-based interventions incorporate not only physical postures but also breathing techniques, mindfulness and relaxation practices. These additional components may provide psychological benefits and contribute to improved quality of life. However, the extent to which yoga-based balance exercises are more effective than conventional training methods in improving functional mobility and overall well-being among elderly individuals remains an important research question. Comparative studies are therefore essential to determine which approach produces better outcomes in terms of balance, mobility and quality of life. Such research can provide valuable evidence for healthcare professionals, physiotherapists and fitness instructors when designing fall-prevention programs for older adults. Furthermore, understanding the relative benefits of yoga and conventional training may help develop integrated exercise programs that combine the strengths of both approaches to promote healthy aging and improve the quality of life among elderly populations.⁵

2. Review of Literature

Studies on Balance Training in Elderly

Balance training has been widely studied as an effective strategy for improving postural stability, functional mobility and reducing fall risk among elderly individuals. Age-related physiological changes such as decreased muscle strength, impaired proprioception and reduced vestibular function contribute to instability and increase the likelihood of falls. These changes make balance training an essential component of geriatric rehabilitation and fall-prevention programs. Research has demonstrated that structured balance training programs significantly improve mobility and stability in older adults. Exercises such as weight shifting, one-leg standing, gait training and strengthening of lower-limb muscles are commonly used in physiotherapy programs. These exercises enhance neuromuscular coordination and help older adults maintain body equilibrium during daily activities. Several studies have reported that balance training can reduce the incidence of falls and improve physical performance in older adults. Programs that challenge balance through controlled movements and functional activities have been found to improve gait speed, reaction time and postural control. Such interventions are especially beneficial for community-dwelling older adults who experience reduced mobility and fear of falling.⁶ Furthermore, fall-prevention exercise programs such as the Otago Exercise Programme and similar physiotherapy-based interventions have shown positive effects on improving strength, balance and functional independence among elderly individuals. These programs focus on lower-limb strengthening and balance exercises designed to reduce fall risk and improve overall mobility. Overall, the literature indicates that balance training plays a crucial role in improving physical functioning, preventing falls and enhancing the independence of older adults. However, researchers have also suggested exploring complementary exercise approaches that may provide additional psychological and holistic health benefits.

Research on Yoga and Fall Prevention

⁴ American College of Sports Medicine. Exercise Guidelines for Older Adults.

⁵ Wayne P.M. Kaptchuk T.J. Challenges inherent to yoga research. *Journal of Alternative and Complementary Medicine*.

⁶ Sherrington C. et al. Exercise interventions for preventing falls in older people.

Yoga has gained increasing attention in recent years as a potential intervention for improving balance and preventing falls among older adults. Yoga involves a combination of physical postures, breathing techniques and relaxation practices that may improve strength, flexibility and postural control. Several studies have demonstrated that yoga practice can significantly improve balance and mobility in elderly populations. For example, a randomized controlled trial evaluating a 12-week Iyengar yoga program found that participants who practiced yoga showed significant improvements in standing balance, walking speed and sit-to-stand performance compared with the control group.⁷ These findings suggest that yoga may serve as an effective exercise intervention for improving mobility and stability among older adults. Recent research has also highlighted the psychological benefits of yoga for elderly individuals. A quasi-experimental study involving older adults aged 65–85 years reported that a structured 12-week yoga program significantly improved balance and functional mobility while also reducing fear of falling, anxiety and depression among participants. The study concluded that yoga can be a safe and effective non-pharmacological approach for promoting healthy aging and enhancing mental well-being. In addition, systematic reviews examining yoga interventions have suggested that yoga may improve fall-related physical functions such as balance, gait performance and lower-limb strength. However, the results across studies remain somewhat inconsistent due to differences in study design, participant characteristics and yoga protocols.⁸ Despite these variations, the overall body of evidence indicates that yoga may play a valuable role in improving balance and reducing fall risk among elderly populations.

Studies Comparing Alternative Therapies with Physiotherapy

In recent years, researchers have explored various alternative or complementary therapies, such as yoga, tai chi and mind–body exercise programs, as potential interventions for improving balance and preventing falls in elderly populations. These therapies often focus on controlled movement, body awareness and relaxation techniques that may enhance both physical and psychological health. Studies comparing alternative exercise approaches with conventional physiotherapy have shown promising results. For example, randomized controlled trials investigating yoga and tai chi programs have found that these interventions can improve balance, reduce pain and enhance quality of life among older adults living in residential care settings.⁹ Although some studies did not observe statistically significant reductions in fall incidence, improvements were reported in mobility, physical functioning and overall well-being. Similarly, mind–body exercise programs emphasize slow and controlled movements, which may improve proprioception, coordination and neuromuscular control. Compared with traditional physiotherapy exercises that primarily focus on strengthening and functional training, alternative therapies often incorporate mental relaxation and breathing techniques, which may contribute to stress reduction and psychological health. Consequently, researchers have suggested that combining physiotherapy-based balance training with alternative exercise approaches may produce more comprehensive health benefits for elderly individuals. Such integrative exercise programs may address both the physical and psychological aspects of aging.

3. Research Gaps

Despite the growing body of research on balance training and yoga interventions for elderly individuals, several gaps remain in the existing literature. First, many studies have focused on evaluating either conventional balance training or yoga interventions separately rather than directly

⁷Tiedemann A. et al. A 12-Week Iyengar Yoga Program Improved Balance and Mobility in Older Community-Dwelling People.

⁸Systematic review on yoga and fall-related physical functions in older adults.

⁹Randomized controlled trial examining yoga and tai chi for balance and falls in residential care settings

comparing their effectiveness. As a result, limited evidence exists regarding which approach produces greater improvements in functional mobility and quality of life among elderly populations. Second, many existing studies have relatively small sample sizes or short intervention durations, which may limit the generalizability of their findings. Longer intervention periods and larger participant groups are needed to better understand the long-term effects of yoga-based exercises and conventional balance training on fall prevention and functional mobility.

Third, while several studies have evaluated physical outcomes such as balance, gait and muscle strength, fewer studies have examined the broader impact of these interventions on psychological well-being and quality of life among elderly individuals. Considering that aging affects both physical and mental health, it is important to assess the holistic benefits of exercise interventions. Finally, there is a need for more experimental studies comparing yoga-based balance exercises with conventional physiotherapy-based balance training programs in a controlled research design. Such comparative studies can provide valuable evidence to guide healthcare professionals in selecting the most effective exercise interventions for improving mobility, preventing falls and enhancing quality of life among older adults.

4. Objectives of the Study

The present study aims to examine and compare the effectiveness of yoga-based balance exercises and conventional balance training in improving functional mobility and quality of life among elderly individuals. With the growing aging population and increasing incidence of falls among older adults, it is essential to identify effective and sustainable exercise interventions that can enhance mobility, maintain independence and promote overall well-being. The specific objectives of the study are described below.

1. To evaluate the effect of yoga-based balance exercises on functional mobility among elderly individuals

One of the primary objectives of this study is to assess the impact of yoga-based balance exercises on the functional mobility of elderly individuals. Functional mobility refers to the ability of a person to move independently and safely while performing daily activities such as walking, standing up from a chair, turning and maintaining balance while moving. Aging is often associated with a decline in muscle strength, coordination and postural stability, which can significantly impair mobility and increase the risk of falls.¹⁰ Yoga-based exercises incorporate various physical postures (asanas), breathing techniques (pranayama) and relaxation practices that may improve balance, flexibility, muscle strength and body awareness. Many yoga postures require controlled body movements and sustained alignment, which can strengthen stabilizing muscles and improve neuromuscular coordination. Through regular practice, these exercises may enhance postural control and increase confidence in performing daily activities. Therefore, this objective seeks to evaluate whether participation in a structured yoga program can significantly improve the functional mobility of elderly individuals.

2. To evaluate the effect of conventional balance training on functional mobility

Another important objective of the study is to examine the effectiveness of conventional balance training programs in improving functional mobility among elderly individuals. Conventional balance training is commonly used in physiotherapy and rehabilitation settings to address mobility limitations

¹⁰Shumway-Cook A. Woollacott M. Motor Control: Translating Research into Clinical Practice.

and reduce the risk of falls. Such training typically includes exercises such as static standing balance, weight-shifting activities, gait training and strengthening exercises targeting the lower limbs.¹¹ These exercises are designed to enhance muscle strength, coordination and stability, which are essential for maintaining balance during both static and dynamic movements. By improving neuromuscular control and reaction time, conventional balance training can help elderly individuals respond more effectively to sudden changes in posture or external disturbances. The purpose of this objective is to determine the extent to which conventional physiotherapy-based balance training improves functional mobility and physical performance among older adults.

3. To compare the impact of yoga-based exercises and conventional balance training on quality of life

In addition to evaluating physical outcomes, this study also aims to compare the effects of yoga-based exercises and conventional balance training on the quality of life of elderly individuals. Quality of life is a multidimensional concept that includes physical health, psychological well-being, social relationships and environmental factors that influence an individual's overall life satisfaction.¹² Elderly individuals often experience physical limitations, emotional stress, social isolation and reduced independence, all of which can negatively affect their quality of life. Exercise interventions have been shown to improve physical health and psychological well-being; however, different types of exercise may influence these aspects in different ways. Yoga, for example, emphasizes not only physical movement but also breathing control, relaxation and mindfulness, which may help reduce stress, anxiety and depression. By comparing the outcomes of yoga-based exercises and conventional balance training, this objective seeks to identify which intervention produces greater improvements in the overall quality of life among elderly participants.

4. To determine which intervention is more effective for improving overall well-being

The final objective of the study is to determine which of the two interventions—yoga-based balance exercises or conventional balance training—is more effective in promoting the overall well-being of elderly individuals. Overall well-being encompasses both physical health and psychological functioning, including improved mobility, reduced fall risk, enhanced mental health and greater independence in daily activities.¹³ While conventional balance training primarily focuses on improving physical stability and functional mobility, yoga provides a holistic approach that addresses physical, mental and emotional aspects of health. By comparing the outcomes of both interventions through standardized assessment tools, this study aims to identify the most beneficial approach for improving the health and well-being of elderly individuals. The findings may provide valuable insights for healthcare professionals, physiotherapists and community health programs seeking effective strategies to support healthy aging and fall prevention.

5. Research Methodology

5.1 Research Design

The study employed a comparative experimental research design to examine the effectiveness of yoga-based balance exercises compared with conventional balance training among elderly individuals. Participants were assessed using pre-test and post-test measurements to evaluate changes in functional

¹¹Sherrington C. et al. Exercise interventions for preventing falls in older people.

¹²World Health Organization. WHOQOL: Measuring Quality of Life.

¹³Lord S.R. Sherrington C. Menz H.B. Falls in Older People: Risk Factors and Strategies for Prevention.

mobility and quality of life following the intervention. The design allowed for systematic comparison between two structured exercise interventions delivered over the same time period.

5.2 Participants

The study included elderly individuals aged 60 years and above residing in community settings. Participants were selected based on the following criteria:

- Inclusion Criteria
- Age 60 years and above
- Ability to walk independently or with minimal assistance
- Willingness to participate in exercise training programs
- Exclusion Criteria
- Severe neurological disorders affecting mobility
- Recent fractures or major surgeries
- Severe cognitive impairment

A total sample of 40 participants was selected and randomly assigned to two groups with 20 participants in each group.

5.3 Ethical Consideration

Ethical approval for the study was obtained from the Institutional Ethics Committee prior to the commencement of data collection. All participants were informed about the purpose and procedures of the study and written informed consent was obtained before participation. Confidentiality and anonymity of participant information were strictly maintained throughout the study.

5.4 Sampling Technique

Participants were selected using a purposive sampling technique ensuring that only elderly individuals who met the specified inclusion criteria were included in the study. After recruitment participants were randomly allocated into two intervention groups using simple random allocation to reduce allocation bias. Each group consisted of 20 participants resulting in a total sample size of 40 individuals.

5.5 Intervention Programs

Yoga-Based Balance Exercise Program

Participants in the yoga group performed structured yoga sessions including:

- Tadasana (Mountain Pose)
- Vrikshasana (Tree Pose)
- Trikonasana (Triangle Pose)
- Bhujangasana (Cobra Pose)
- Pranayama breathing exercises
- Relaxation techniques

The intervention programs were conducted for a fixed duration of 8 weeks with three sessions per week and each session lasting approximately 45 minutes under the supervision of trained instructors.

Yoga Posture	Duration	Repetitions
Tadasana (Mountain Pose)	20 seconds	3 repetitions
Vrikshasana (Tree Pose)	15 seconds each side	3 repetitions

Trikonasana (Triangle Pose)	15 seconds each side	3 repetitions
Bhujangasana (Cobra Pose)	20 seconds	3 repetitions
Pranayama Breathing	5 minutes	Continuous
Relaxation (Shavasana)	5 minutes	End of session

The yoga intervention protocol was standardized to ensure consistency across sessions. Each posture was performed for a specified duration with controlled breathing and proper body alignment. Sessions were conducted under the supervision of a trained yoga instructor to ensure safety and correct technique.

5.6 Conventional Balance Training Program

Participants in the conventional training group performed physiotherapy-based exercises such as:

- Static standing balance exercises
- Weight-shifting activities
- Gait training
- Heel-to-toe walking
- Lower limb strengthening exercises

Exercise	Sets	Repetitions
Static Standing Balance	3	20 seconds hold
Weight Shifting	3	15 repetitions
Heel-to-Toe Walking	3	10 steps
Gait Training	3	10 meters
Lower Limb Strengthening	3	12 repetitions

The conventional balance training program was structured to ensure uniform exercise dosage across participants. All exercises were supervised by a physiotherapist and rest intervals of approximately 1 minute between sets were provided to avoid fatigue.”

5.7 Outcome Measures

To evaluate the effectiveness of the interventions, the following standardized assessment tools were used:

1. Functional Mobility

- Timed Up and Go (TUG) Test
- Gait speed test

2. Quality of Life

WHOQOL-BREF questionnaire

These assessments were conducted before and after the intervention period.

5.8 Data Collection Procedure

Data were collected in three stages:

- Pre-test assessment of participants' functional mobility and quality of life.
- Implementation of intervention programs for both groups over a period of 8–12 weeks.

- Post-test assessment using the same evaluation tools to measure improvements.

5.9 Data Analysis

The collected data were analyzed using Statistical Package for Social Sciences (SPSS) version XX. Descriptive statistics including mean and standard deviation were calculated for all outcome variables. The Shapiro–Wilk test was used to examine the normality of the data distribution. To evaluate within-group improvements a paired sample t-test was conducted to compare pre-test and post-test scores. An independent sample t-test was used to compare differences between the yoga and conventional training groups. Statistical significance was set at $p < 0.05$ and 95% confidence intervals were calculated for major outcome measures.

6. Results

The results indicated improvements in functional mobility and quality of life in both intervention groups following the 8-week training program. Table 1 presents the pre-test and post-test scores for the Timed Up and Go (TUG) test.

Group	Pre-test TUG (Mean±SD)	Post-test TUG (Mean±SD)	p-value
Yoga Group	12.4 ± 2.1 sec	9.1 ± 1.8 sec	0.001
Conventional Training	12.7 ± 2.3 sec	10.8 ± 2.0 sec	0.012

Participants in both groups demonstrated statistically significant improvements in mobility; however the yoga group showed a greater reduction in TUG scores indicating improved functional mobility.

7. Discussion

The findings of the present study suggest that both yoga-based balance exercises and conventional balance training are effective interventions for improving functional mobility among elderly individuals. These findings are consistent with previous research indicating that structured exercise programs can enhance postural stability and reduce fall risk in older adults.¹⁴ However, the yoga-based intervention demonstrated greater improvements in overall quality of life compared with conventional balance training. This may be attributed to the holistic nature of yoga, which integrates physical activity with breathing exercises and relaxation techniques. Such practices may contribute to improved psychological well-being and reduced stress levels.¹⁵ Furthermore, yoga postures require controlled body movements and sustained balance, which may strengthen stabilizing muscles and enhance neuromuscular coordination. These factors may explain the greater improvements in functional mobility observed in the yoga group. The results also support the growing body of evidence suggesting that mind–body exercise programs such as yoga may provide comprehensive benefits for elderly populations. Integrating yoga into fall-prevention programs may therefore represent an effective strategy for promoting healthy aging.

8. Limitations of the Study

While the present study provides valuable insights into the effectiveness of yoga-based balance exercises and conventional balance training among elderly individuals, several limitations should be acknowledged. Recognizing these limitations helps in understanding the scope of the findings and provides direction for future research.

¹⁴Sherrington C. et al. Exercise for preventing falls in older people.

¹⁵Woodyard C. Exploring the therapeutic effects of yoga and its ability to increase quality of life. *International Journal of Yoga*.

1. Limited Sample Size

One of the primary limitations of the study is the relatively small sample size. The number of participants involved in the study was limited, which may affect the generalizability of the findings to the larger elderly population. A larger sample size would allow for more accurate statistical analysis and provide more reliable conclusions regarding the effectiveness of the interventions. Future studies should consider involving a greater number of participants from different demographic backgrounds to strengthen the validity of the results.

2. Short Duration of Intervention

Another limitation of the study is the relatively short duration of the intervention program. The exercise interventions were conducted over a limited period, which may not fully reflect the long-term benefits of yoga-based balance exercises or conventional balance training. Long-term practice of these exercises could produce more significant improvements in mobility, balance and quality of life. Therefore, future research should include longer intervention periods and follow-up assessments to examine the sustained effects of these training programs.

3. Limited Geographic Area

The participants in this study were selected from a specific geographic area or community. As a result, the findings may not be representative of elderly populations from different regions, cultures or socioeconomic backgrounds. Environmental factors, lifestyle habits and access to healthcare services may vary across different populations and could influence the outcomes of exercise interventions. Future research should include participants from multiple regions to enhance the generalizability of the findings.

4. Variability in Participant Health Conditions

Elderly individuals often have varying levels of physical fitness, health conditions and mobility limitations. Some participants may have underlying medical conditions such as arthritis, hypertension or diabetes, which could influence their ability to perform exercises and affect the outcomes of the study. Although efforts were made to select participants based on specific inclusion criteria, individual differences in health status may have influenced the results.

5. Dependence on Self-Reported Measures

The study included the assessment of quality of life through questionnaires and self-reported responses. Such measures may be subject to personal bias, differences in interpretation or social desirability effects. Participants may sometimes provide responses that reflect what they believe is expected rather than their actual experiences. Although standardized questionnaires were used, objective measures combined with self-reported assessments could provide a more comprehensive evaluation.

6. Lack of Long-Term Follow-Up

The study primarily focused on evaluating the immediate effects of the intervention programs after the completion of the training period. However, it did not assess whether the improvements in mobility and quality of life were maintained over time. Long-term follow-up studies would help determine whether the benefits of yoga-based exercises and conventional balance training persist after the intervention ends.

7. Recommendations

Based on the findings of the study on the effectiveness of yoga-based balance exercises and conventional balance training among elderly individuals, several recommendations can be made for healthcare professionals, community organizations and future researchers. These recommendations aim to enhance fall prevention strategies, improve functional mobility and promote overall well-being among older adults.

1. Incorporation of Yoga-Based Balance Exercises in Geriatric Health Programs

The results of the study suggest that yoga-based balance exercises can significantly improve functional mobility and enhance the quality of life among elderly individuals. Therefore, it is recommended that healthcare institutions, rehabilitation centers and community health programs incorporate structured yoga sessions as part of their geriatric care and wellness programs. Yoga offers a holistic approach that not only improves physical strength and balance but also promotes mental relaxation and emotional well-being. Regular practice of yoga can help elderly individuals maintain independence, reduce the risk of falls and improve their overall health status.

2. Integration of Yoga and Conventional Balance Training

Although yoga-based exercises demonstrated greater benefits in certain aspects, conventional balance training also plays an important role in improving muscle strength and postural stability. Therefore, it is recommended that healthcare professionals consider integrating both approaches in exercise programs for elderly individuals. A combined program that includes physiotherapy-based exercises along with yoga postures and breathing techniques may provide comprehensive benefits by addressing both physical and psychological aspects of health. Such integrative programs could be particularly effective in fall prevention and rehabilitation settings.

3. Promotion of Community-Based Exercise Programs for the Elderly

Community centers, senior citizen organizations and public health agencies should promote accessible exercise programs specifically designed for elderly populations. Regular physical activity programs that include yoga and balance training can help older adults stay active, socially engaged and physically independent. These programs should be supervised by trained professionals to ensure safety and proper technique. Community-based initiatives can also raise awareness about the importance of maintaining balance and mobility as part of healthy aging.

4. Training of Healthcare Professionals and Instructors

Healthcare professionals, physiotherapists and fitness instructors who work with elderly populations should receive specialized training in both yoga-based interventions and conventional balance training techniques. Proper knowledge and training will enable instructors to design safe and effective exercise programs tailored to the specific needs and limitations of elderly individuals. Training programs should emphasize safe practice, appropriate modifications of exercises and monitoring of participants to prevent injuries.

5. Early Implementation of Fall-Prevention Programs

Fall prevention strategies should be introduced at an early stage in the aging process. It is recommended that individuals approaching older adulthood begin engaging in regular balance training and yoga exercises before significant mobility decline occurs. Early implementation of such programs may help maintain strength, flexibility and balance, thereby reducing the likelihood of falls and associated injuries later in life.

6. Encouragement of Regular Physical Activity among Elderly Individuals

Elderly individuals should be encouraged to participate in regular physical activity to maintain their physical and mental health. Family members, caregivers and healthcare providers play an important role in motivating older adults to engage in safe and enjoyable exercise routines. Programs that combine social interaction with physical activity may increase adherence and make exercise more appealing for elderly participants.

7. Recommendations for Future Research

Future research should consider expanding the scope of the present study by including larger sample sizes and longer intervention periods. Studies involving diverse populations and different geographical locations would provide more generalizable results. Additionally, future research could explore the long-term effects of yoga-based balance exercises on fall prevention, mental health and chronic disease management among elderly individuals. Researchers may also investigate the effectiveness of different styles of yoga or combined exercise interventions in improving functional mobility and quality of life.

8. Policy-Level Support for Elderly Fitness Programs

Government agencies and public health organizations should support the development of policies that promote physical activity among elderly populations. Funding and infrastructure should be provided to establish accessible exercise facilities, training programs and community health initiatives that encourage active aging. Such policies would contribute to reducing healthcare costs associated with fall-related injuries and improving the overall quality of life among older adults.

Conclusion

The present study aimed to examine the effectiveness of yoga-based balance exercises in comparison with conventional balance training on functional mobility and quality of life among elderly individuals. Aging is often associated with a gradual decline in physical abilities such as muscle strength, coordination, flexibility and postural stability. These age-related changes significantly increase the risk of falls and reduce the ability of older adults to perform daily activities independently. As a result, identifying effective exercise interventions that can improve balance and mobility is essential for promoting healthy aging and enhancing the quality of life among elderly populations. The findings of this study indicate that both yoga-based balance exercises and conventional balance training are beneficial in improving functional mobility among elderly individuals. Participants in both groups showed improvements in mobility, balance control and the ability to perform daily activities more safely and efficiently after completing the intervention program. These results support previous research suggesting that structured exercise programs play a crucial role in reducing fall risk and improving physical functioning in older adults. Regular participation in balance training exercises helps strengthen lower limb muscles, improve coordination and enhance neuromuscular control, which are essential for maintaining postural stability.

However, the results also suggest that yoga-based balance exercises may provide additional benefits beyond those achieved through conventional balance training alone. Yoga integrates physical postures, breathing techniques and relaxation practices, which together promote a holistic approach to health and well-being. The study findings showed that elderly participants practicing yoga not only improved their functional mobility but also reported better psychological well-being and overall quality of life. This may be attributed to the calming and stress-reducing effects of yoga, which can help reduce anxiety, enhance mental clarity and improve emotional balance. Another important aspect highlighted by the study is the positive impact of exercise on the quality of life of elderly individuals. Quality of life is influenced by several factors, including physical health, psychological well-being, social

33 participation and independence in daily living. The results suggest that exercise programs, particularly those incorporating yoga, can significantly enhance these aspects of life by promoting physical fitness, reducing fear of falling and encouraging active participation in daily activities. Elderly individuals who maintain good mobility and balance are more likely to remain independent and socially engaged, which contributes to a greater sense of satisfaction and well-being.

38
10
13
16
45
1
2
Despite these positive findings, it is important to recognize that conventional balance training also plays a vital role in improving physical stability and functional performance. Physiotherapy-based exercises are specifically designed to target muscle strength, coordination and balance control. Therefore, both interventions can be considered effective approaches for fall prevention and mobility enhancement. The choice of intervention may depend on individual preferences, health conditions and accessibility to training programs. Overall, the findings of this study highlight the importance of incorporating regular physical activity into the daily routine of elderly individuals. Exercise programs that focus on balance and mobility can significantly reduce the risk of falls, improve independence and enhance overall quality of life. In particular, yoga-based balance exercises appear to offer a comprehensive approach by addressing both physical and psychological aspects of health. In conclusion, the study suggests that yoga-based balance exercises can serve as an effective alternative or complementary approach to conventional balance training for elderly individuals. Integrating yoga into geriatric health programs and rehabilitation settings may provide significant benefits in terms of improved mobility, reduced fall risk and enhanced well-being. Encouraging older adults to engage in safe and structured exercise programs will play an important role in promoting healthy aging and supporting a higher quality of life in later years.



SELF PUBLISHED AUTHOR'S PUBLICATION